

On compression of Bruhat–Tits buildings

YURII A. NERETIN

Consider an affine Bruhat–Tits building Lat_n of the type A_{n-1} and the complex distance in Lat_n , i.e., the complete system of invariants of a pair of vertices of the building. An elements of the Nazarov semigroup is a lattice in the duplicated p -adic space $\mathbb{Q}_p^n \oplus \mathbb{Q}_p^n$. We investigate behavior of the complex distance with respect to the natural action of the Nazarov semigroup on the building.

It is well known that affine Bruhat–Tits buildings (see, for instance, [1], [3], [7]) are right p -adic analogs of Riemannian noncompact symmetric spaces. This analogy has no a priori explanations, but it exists on the levels of geometry, harmonic analysis, and special functions. Some known examples of this parallel are

- Gindikin–Karpelevich’s and Macdonald’s Plancherel formulae, see [7];
- existing of a general theory including Hall–Littlewood polynomials and zonal spherical functions, see [8];
- similarity in the geometries at infinity (Semple–Satake type boundaries, Martin boundary etc.), see [4], [13];
- parallel between Horn–Klyachko (see [2]) triangle inequality for eigenvalues of Hermitian matrices¹ and the triangle inequality for complex distance in buildings²
- Existence of a precise parallels of the harmonic analysis of the Berezin kernels and of the matrix beta-functions on the level of buildings³, see [16].

In this note, we continue this parallel and obtain a (quite simple) analog of the compression of angles theorem (this theorem appeared in [11] as a lemma for boundedness of some integral operators in Fock spaces, see its proof in [12], 6.3, and some geometrical discussion in [14], see also [5] on its extension to exceptional groups) for buildings of the type A_n .

1. Formulation of result

1.1. Notation. We denote by $\mathbb{Q}_p$ the p -adic field. By $|\cdot|$, we denote the norm on $\mathbb{Q}_p$. Let $\mathbb{O}_p$ be the ring of p -adic integers. Recall that $\mathbb{O}_p$ consists of elements with norm ≤ 1 . By $\mathbb{Q}_p^n$ we denote the coordinate n -dimensional linear space over $\mathbb{Q}_p$.

By $\text{GL}_n(\mathbb{Q}_p)$ we denote the group of invertible $n \times n$ matrices over $\mathbb{Q}_p$. By $\text{GL}_n(\mathbb{O}_p)$ we denote the group of matrices $g \in \text{GL}_n(\mathbb{Q}_p)$ such that all the matrix elements of g and g^{-1} are integer.

¹Interpretation of this inequality in terms of geometry of symmetric spaces is discussed in [15]

²Let $R, S, T \in \text{Lat}_n$ (see notations below). To describe possible collections of $3n$ integers $\{k_j(R, S)\}, \{k_j(R, T)\}, \{k_j(S, T)\}$. It can be easily reduced to the following (unexpectedly nontrivial) question solved in 1960-s by Ph.Hall and T.Klein. Let A, B be finite Abelian p -groups. For which Abelian p -groups C there exists a subgroup A' isomorphic to A such that $C/A' \simeq B$, for a solution, see [8], II.3.

³The matrix gamma-function corresponds to the zeta-function of Tamagawa, see [8], V.4.

1.2. Space of lattices. A lattice in $\mathbb{Q}_p^n$ is a compact open $\mathbb{O}_p$ -submodule. Recall, that any lattice R can be represented in the form

$$R = \mathbb{O}f_1 \oplus \cdots \oplus \mathbb{O}f_n, \quad (0.1)$$

where $f_1, \dots, f_n$ is some basis in $\mathbb{Q}_p^n$, see [18], §II.2. We denote by Lat_n the space of lattices in $\mathbb{Q}_p^n$ (this space is countable, and a natural topology on Lat_n is discrete).

The group $\text{GL}_n(\mathbb{Q}_p)$ acts transitively on Lat_n . The stabilizer of the lattice $\mathbb{O}_p^n$ is $\text{GL}_n(\mathbb{O})$. Thus,

$$\text{Lat}_n = \text{GL}_n(\mathbb{Q}_p)/\text{GL}_n(\mathbb{O}_p).$$

1.3. Complex distance.

THEOREM (see, for instance, [18], Theorem 2.2.2). *Let R, S be two lattices in $\mathbb{Q}_p^n$.*

a) *There exists a basis $f_1, \dots, f_n \in \mathbb{Q}_p^n$ such that*

$$R = \sum_{j=1}^n \mathbb{O}_p f_j, \quad S = \sum_{j=1}^n p^{-k_j(R,S)} \mathbb{O}_p f_j \quad (0.2)$$

Integer numbers

$$k_1(R, S) \geq k_2(R, S) \geq \dots \geq k_n(R, S)$$

are uniquely determined by the lattices R, S .

b) *If S, R and S', R' be elements of Lat_n satisfying*

$$k_j(R, S) = k_j(R', S')$$

then there exists $g \in \text{GL}_n(\mathbb{Q}_p)$ such that

$$R = g \cdot R', \quad S = g \cdot S'$$

We say that $k_j(R, S)$ is the *complex distance* between S, R (this structure is the analog of the complex distance or stationary angles in symmetric spaces, see, for instance, [15]).

1.4. Product of relations. Let V, W be sets. A relation $H : V \rightrightarrows W$ is a subset in $V \times W$.

Let $H : V \rightrightarrows W, G : W \rightrightarrows Y$ be relations. Their product $G \cdot H : V \rightrightarrows Y$ is a relation consisting of all pairs $v \times y \in V \times Y$ for which there exists $w \in W$ satisfying $v \times w \in H, w \times y \in G$.

1.5. Nazarov semigroup. An element of the Nazarov semigroup Γ_n is a lattice in $\mathbb{Q}_p^n \times \mathbb{Q}_p^n$. The product in the Nazarov semigroup is the product of relations.

REMARK. It is possible to consider elements of the group $\text{GL}_n(\mathbb{Q}_p)$ as points of the Nazarov semigroup at infinity. Indeed, for $g \in \text{GL}_n(\mathbb{Q}_p)$ consider a sequence of lattices $Z_j \subset \mathbb{Q}_p^n \oplus \mathbb{Q}_p^n$ such that

$$\bigcap_{j=1}^{\infty} \bigcup_{k=j}^{\infty} Z_j$$

coincides with the graph of g . For each lattice $R \subset \mathbb{Q}_p^n$, we have $Z_j \cdot R = g \cdot R$ starting from some place j .

REMARK. Nazarov semigroups are p -adic analogues of semigroups of linear relations (see [12]). The main nontrivial property of the Nazarov semigroups is the following statement. The Weil representation of $\mathrm{Sp}(2n, \mathbb{Q}_p)$ (see [17]) admits a canonical extension to representation of a Nazarov semigroup related to buildings of the type C_n , see [10], [9]. It seems that this statement must have many corollaries (due to possibilities of the Howe duality), these corollaries never were discussed in literature.

1.6. Action of the Nazarov semigroup on Lat_n . Consider $H \in \Gamma_n$, $R \in \mathrm{Lat}_n$. We define $HR \in \mathrm{Lat}_n$ as set of all $w \in \mathbb{Q}_p^n$ such that there exists $v \in R$ satisfying $v \times w \in H$.

1.7. Result of the paper. The main result of this paper is the following theorem.

THEOREM. *Let H be an element of the Nazarov semigroup Γ_n . Let $R, S \in \mathrm{Lat}_n$. Then for each j the number $k_j(H \cdot R, H \cdot S)$ lies between 0 and $k_j(R, S)$, i.e.,*

$$\begin{aligned} k_j(R, S) \geq 0 & \quad \text{implies} \quad 0 \leq k_j(H \cdot R, H \cdot S) \leq k_j(R, S) \\ k_j(R, S) \leq 0 & \quad \text{implies} \quad 0 \geq k_j(H \cdot R, H \cdot S) \geq k_j(R, S) \end{aligned}$$

REMARK. In the case of Riemannian symmetric spaces, there exists an inverse theorem about characterization of all angle-compressive maps [14]. I do not know such statements in the p -adic case.

2. Proof

2.1. Norms. A *norm* on $\mathbb{Q}_p^n$ is a function $v \mapsto \|v\|$ from $\mathbb{Q}_p^n$ to the set

$$\{\dots, p^{-1}, 1, p, p^2, \dots\}$$

satisfying the conditions

1. $\|v + w\| \leq \max(\|v\|, \|w\|)$
2. $\|\lambda v\| = |\lambda| \|v\|$ for $\lambda \in \mathbb{Q}_p$, $v \in \mathbb{Q}_p^n$.

The set of all v , such that $\|v\| \leq 1$, is a lattice.

Conversely, let L be a lattice. We define the corresponding norm $\|\cdot\|_L$ by the rule

$$\|v\|_L := \left(\min_{k: p^k v \in L} p^k \right)^{-1}$$

2.2. Minimax characterization of the complex distance. The following statement is an exact analogue of the standard minimax characterization of the eigenvalues (see, for instance, [6]) and of the stationary angles (see, for instance [15]). The proofs also is similar.

LEMMA 1. *For $L, M \in \mathrm{Lat}_n$*

$$\begin{aligned} \text{a) } p^{k_j(R,S)} &= \max_{W: \dim W=j, W \subset \mathbb{Q}_p^n} \left\{ \min_{v \in W, v \neq 0} \frac{\|v\|_S}{\|v\|_R} \right\} \\ \text{b) } p^{k_j(R,S)} &= \min_{W: W \subset \mathbb{Q}_p^n, \text{codim } W=j-1} \left\{ \max_{v \in W, v \neq 0} \frac{\|v\|_S}{\|v\|_R} \right\} \end{aligned}$$

PROOF. Let R, S have the form (0.2). Consider the subspace $Y = \mathbb{Q}_p f_j \oplus \dots \oplus \mathbb{Q}_p f_n$. We have $\|v\|_R / \|v\|_S \leq p^{k_j}$ for each $v \in Y$.

If $\dim W_j = j$, then $Y \cap W \neq 0$, and hence $\min_{v \in W} \|v\|_R / \|v\|_S \leq p^{k_j}$.

For $W = \mathbb{Q}_p f_1 \oplus \dots \oplus \mathbb{Q}_p f_j$ we have $\min_{v \in W} \|v\|_R / \|v\|_S = p^{k_j}$. $\square$

2.3. Another characterization of the complex distance. The following statement is obvious.

LEMMA 2. *Let $L, M \in \text{Lat}_n$. Then we have the following isomorphisms of finite Abelian p -groups:*

$$\begin{aligned} M/(L \cap M) &= \bigoplus_{j: k_j(L,M) > 0} \mathbb{Z}/p^{k_j(L,M)} \mathbb{Z} \\ L/(L \cap M) &= \bigoplus_{j: k_j(L,M) < 0} \mathbb{Z}/p^{-k_j(L,M)} \mathbb{Z} \end{aligned}$$

2.4. Duality and complex distance. Denote by $(\mathbb{Q}_p^n)^\square$ the linear space dual to $\mathbb{Q}_p^n$. For a lattice $L \in \text{Lat}_n$ we consider the *dual lattice* $L^\square \subset (\mathbb{Q}_p^n)^\square$ consisting of linear functionals $f \in (\mathbb{Q}_p^n)^\square$ such that

$$v \in L \implies f(v) \in \mathbb{O}_p$$

LEMMA 3. $k_j(L, M) = k_j(M^\square, L^\square)$.

2.5. Complex distance and intersections of lattices.

LEMMA 4. *Let $L, M, N \in \text{Lat}_n$. Then for all j , the number $k_j(L \cap M, L \cap N)$ lies between 0 and $k_j(M, N)$.*

This statement is a corollary of Lemma 1 and the following Lemma 5.

LEMMA 5. *For $v \in \mathbb{Q}_p^n$, the number $\|v\|_{L \cap M} / \|v\|_{L \cap N}$ lies between 1 and $\|v\|_M / \|v\|_N$.*

PROOF. Denote

$$\log_p \|v\|_L = l, \quad \log_p \|v\|_M = m, \quad \log_p \|v\|_N = z,$$

Then

$$\log_p \|v\|_{L \cap M} = \max(l, m), \quad \log_p \|v\|_{L \cap N} = \max(l, z)$$

It remains to observe that $\max(l, m) - \max(l, z)$ lies between 0 and $l - z$.

2.6. Complex distance and sums of lattices.

LEMMA 6. *Let $L, M, N \in \text{Lat}_n$. Then $k_j(L + M, L + N)$ lies between 0 and $k_j(M, N)$.*

PROOF. For $P, Q \in \text{Lat}_n$, we have

$$(P + Q)^\square = P^\square \cap Q^\square$$

By the duality consideration, our lemma is equivalent to Lemma 4.

2.7. Notation. Let $S \subset \mathbb{Q}_p^n \oplus \mathbb{Q}_p^n$ be an element of the Nazarov semigroup. We define

- its *kernel* $\text{Ker } S = S \cap (\mathbb{Q}_p^n \oplus 0)$,
- the *domain* $\text{Dom } S$ be the image of S under the projection $\mathbb{Q}_p^n \oplus \mathbb{Q}_p^n \rightarrow \mathbb{Q}_p^n \oplus 0$,
- the *image* $\text{Im } S$ be the image of S under the projection $\mathbb{Q}_p^n \oplus \mathbb{Q}_p^n \rightarrow 0 \oplus \mathbb{Q}_p^n$,
- the *indefiniteness* $\text{Indef } S$ be the intersection $S \cap 0 \oplus \mathbb{Q}_p^n$

Evidently, all the sets $\text{Dom } S$, $\text{Ker } S$, $\text{Indef } S$, $\text{Im } S$ are lattices in $\mathbb{Q}_p^n$.

2.8. Proof of Theorem. Let $S \in \Gamma_n$. We have

$$S \subset \text{Dom } S \oplus \text{Im } S, \quad S \supset \text{Ker } S \oplus \text{Indef } S.$$

Consider the (finite) factor-group

$$Y := (\text{Dom } S / \text{Ker } S) \oplus (\text{Im } S / \text{Indef } S)$$

and the image σ of S in Y . It can be easily be checked that σ is a graph of an isomorphism

$$\text{Dom } S / \text{Ker } S \rightarrow \text{Im } S / \text{Indef } S$$

In particular, these two groups are isomorphic.

Hence, by Lemma 2,

$$k_j(\text{Dom } S, \text{Ker } S) = k_j(\text{Im } S, \text{Indef } S)$$

This implies an existence of a (non canonical) element $g_S \in \text{GL}_n(\mathbb{Q}_p)$ such that

$$g_S \cdot \text{Dom } S = \text{Im } S, \quad g_S \cdot \text{Ker } S = \text{Indef } S$$

For any $L \in \text{Lat}_n$, we have

$$S \cdot L = g_S \cdot (L \cap \text{Dom } S) + \text{Indef } S$$

It remains to apply Lemma 4 and 6.

References

- [1] Brown, K., *Buildings*, Springer, 1989.
- [2] Fulton, W., *Eigenvalues of sums of Hermitian matrices (after A. Klyachko)*. Asterisque No. 252 (1998), Exp. No. 845, 5, 255–269.

- [3] Garret, P. *Buildings and classical groups*, 1996
- [4] Guivarch, Y., Ji, L., Taylor, J. *Compactifications of symmetric spaces*. Birkhauser Boston, Inc., Boston, MA, 1998.
- [5] Koufany, Kh., *Contractions of angles in symmetric cones*. Publ. Res. Inst. Math. Sci. 38 (2002), no. 2, 227–243.
- [6] Lidskii, V.B. *Inequalities for eigenvalues and singular values*, addendum to Gantmaher F.R. *Theory of matrices*, Second (1966), Third (1976), Forth (1988) Russian editions.
- [7] Macdonald, I.G. *Spherical functions on a group of p -adic type*, Publ.Ramanujan Inst., Madras, 1972
- [8] Macdonald, I.G. *Symmetric functions and Hall polynomials*, 2-nd ed., Clarendon Press, Oxford, 1995.
- [9] Nazarov, M.L. *The oscillator semigroup over nonarchimedean field*. J. Funct. Anal., 128 (1995), 384–438
- [10] Nazarov, V., Yu. Neretin, Yu., G. Olshanskii, G., *Semi-groupes engendrés par la représentation de Weil du groupe symplectique de dimension infinie*, C. R. Acad. Sci. Paris Sér. I Math. **309**(1989), no. 7, 443–446.
- [11] Neretin, Yu., *On a semigroup of operators in the boson Fock space*, Funct. Anal. Appl. **24**(1990), no. 2, 135–144.
- [12] Neretin Yu., *Categories of symmetries and infinite-dimensional groups*, London Math. Soc. Monographs, N. S., **16**, Oxford University Press, 1996.
- [13] Neretin, Yu. A. *Hinges and the Study-Semple-Satake-Furstenberg-De Concini-Procesi-Oshima boundary*. Kirillov’s seminar on representation theory, 165–230, Amer. Math. Soc. Transl. Ser. 2, 181, Amer. Math. Soc., Providence, RI, 1998.
- [14] Neretin, Yu. A. *Conformal geometry of symmetric spaces, and generalized linear-fractional Kreĭn-Shmulian mappings*. Mat. Sb. 190 (1999), no. 2, 93–122; transl. in Sb. Math. 190 (1999), no. 1-2, 255–283
- [15] Neretin, Yu.A. *Jordan angles and triangle inequality in Grassmannian manifold*, Geometria Dedicata, 86 (2001), 403–432
- [16] Neretin, Yu.A. *Beta-function of Bruhat–Tits buildings and deformation of L^2 on the space of lattices*. Mat. Sbornik, 2003; preprint version is available via <http://xxx.arxiv.org/abs/math/0306079>.
- [17] Weil, A. *Sur certains groupes d’opérateurs unitaires*, Acta Math., 111 (1964), 143–211
- [18] Weil, A. *Basic number theory*, Springer, 1967;

Math. Phys. Group,
Institute for Theoretical and Experimental Physics,
B. Cheremushkinskaya, 25, Moscow -- 117259, Russia
&
University of Vienna
neretin@mccme.ru,